

# The effects of rest and treatment following sport-related concussion: a systematic review of the literature

Kathryn J Schneider,<sup>1</sup> Grant L Iverson,<sup>2</sup> Carolyn A Emery,<sup>1,3,4</sup> Paul McCrory,<sup>5</sup> Stanley A Herring,<sup>6,7</sup> Willem H Meeuwisse<sup>1,3</sup>

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2013-092190>)

<sup>1</sup>Faculty of Kinesiology, Sport Injury Prevention Research Centre, University of Calgary, Calgary, Alberta, Canada

<sup>2</sup>Department of Psychiatry, Faculty of Medicine, University of British Columbia, Vancouver, British Columbia, Canada

<sup>3</sup>Department of Community Health Sciences, Faculty of Medicine, University of Calgary, Calgary, Alberta, Canada

<sup>4</sup>Alberta Children's Hospital Research Institute for Child & Maternal Health, Faculty of Medicine, University of Calgary, Calgary, Alberta, Canada

<sup>5</sup>The Florey Institute of Neuroscience and Mental Health, University of Melbourne & Melbourne Brain Centre, Heidelberg, Victoria, Australia

<sup>6</sup>Departments of Rehabilitation Medicine, Orthopaedics and Sports Medicine and Neurological Surgery, University of Washington, Seattle, Washington, USA

<sup>7</sup>Seattle Sports Concussion Program, Team Physician Seattle Seahawks and Seattle Mariners, Seattle, Washington, USA

## Correspondence to

Dr Kathryn J Schneider, Faculty of Kinesiology, Sport Injury Prevention Research Centre, University of Calgary, 2500 University Drive N.W., Calgary, Alberta, Canada T2N 1N4; [kjschnei@ucalgary.ca](mailto:kjschnei@ucalgary.ca)

Received 15 January 2013  
Accepted 20 January 2013

## ABSTRACT

**Objective** To evaluate the evidence for rest, treatment, and rehabilitation following sport-related concussion (SRC).

**Data sources** PubMed, CINAHL, PsychInfo, Cochrane Controlled Trials Registers, Health STAR, Sport Discus, EMBASE, Web of Science, and ProQuest.

**Study selection** Articles were included if they met the following criteria: original research, reported SRC as a source of injury, and evaluated the effect of rest or treatment.

**Data extraction** Study design, participants, treatment, outcome measures, and key findings.

**Data synthesis** Three studies met the inclusion criteria for evaluating the effects of rest and twelve for treatment. Low-intensity aerobic exercise may be of benefit.

**Conclusions** The current evidence evaluating the effect of rest and treatment following SRC is sparse. An initial period of rest may be of benefit. Low-level exercise and multimodal physiotherapy may be of benefit for those who are slow to recover. There is a strong need for high level studies evaluating the effects of rest and treatment following SRC.

## INTRODUCTION

The current recommendation for managing a sport-related concussion is to rest until symptoms resolve and then follow a protocol of graded exertion prior to returning to sport.<sup>1</sup> These recommendations have been developed from a series of agreement and consensus statements by a panels of experts in the area of sport-related concussion.<sup>1,2</sup> As of the 2008 consensus meeting on concussion in sport, there was a paucity of evidence evaluating the effects of various treatments following a concussion.

The purpose of this systematic review was to evaluate the evidence for management of concussion. The following questions were addressed: (1) When is rest useful and/or beneficial as part of the management recommendations for sport-related concussion? and (2) What treatment and rehabilitation strategies are beneficial following a sport-related concussion?

## METHODS

A literature search was conducted for each research question using the following databases: PubMed, CINAHL, PsychInfo, Cochrane Controlled Trials Registers, Health STAR, Sport Discus, EMBASE, Web of Science and ProQuest. To search for articles evaluating the effect of the rest, the following search

terms were used: (Brain Concussion [MeSH] or Post-Concussion Syndrome [MeSH] or Sport-related concussion [tw] or mild traumatic brain injury [tw]) AND (Rest [MeSH] or Treatment Outcome [MeSH] or Cognitive rest [tw] or Physical exertion [MeSH] or Therapy or Rehabilitation or Treatment) AND Sport [tw]. To search for articles evaluating the effect of treatment, the following search terms were used: (Brain Concussion [MeSH] or Post-Concussion Syndrome [MeSH] or Sport-related concussion [tw] or mild traumatic brain injury [tw]) AND (Therapy [MeSH] or Treatment Outcome [MeSH] or Rehabilitation [MeSH] or treatment [tw] or Exercise therapy [MeSH]) AND (Exercise [MeSH] or Brain training [tw] or Cervical spine [tw] or Neck [MeSH] or Vertigo [MeSH] or Dizziness [MeSH] or Headache [MeSH] or Postural Balance [MeSH] or Cognitive therapy [MeSH] or pharmacotherapy [tw]). Inclusion criteria were determined a priori as follows: (1) report of original research (including randomised controlled trials, cohort studies, quasi-experimental designs, case series, case crossovers and case studies), (2) report of a sport-related concussion as a source of symptoms and (3) evaluation of the effect of a resting period or treatment. In order to include the most up-to-date research, peer-reviewed published articles and abstracts were included. All searches were performed by one investigator (KJS).

## Data extraction

For all included articles, the following data were extracted: study design, participants (sample size, age and gender), treatment (type, intensity and duration), key findings (including effect size and means with 95% CIs which were calculated, if possible, if not presented in the original study) and relevant comments. Each study was graded for level of evidence according to the Oxford Centre for Evidence Based Medicine.<sup>3</sup> Downs and Black criteria were used to assess the quality of the evidence.

## RESULTS

The initial search revealed 749 potential articles evaluating the effects of rest and 1175 articles evaluating the effects of treatment. The initial search results revealed: (i) 6 potential studies evaluating rest, 2 of which met the inclusion criteria for the effect of rest following a sport-related concussion and (ii) 15 abstracts that evaluated the effects of treatment, 10 of which met the inclusion criteria. One additional treatment article was identified and included using a Web of Science citation search,<sup>4</sup> and two others were identified by the

**To cite:** Schneider KJ, Iverson GL, Emery CA, et al. *Br J Sports Med* 2013;**47**:304–307.

authors.<sup>5 6</sup> The study design, study sample, outcome measures, description of rest/intervention, key findings and level of evidence are presented in online supplementary table S1.

### Rest

Two papers met the inclusion criteria for evaluating prescribed rest following a sport-related concussion.<sup>7 8</sup> One uncontrolled retrospective observational study of rest and activity level was identified.<sup>9</sup> The majority of abstracts that were initially screened and were subsequently excluded did not specifically evaluate the effects of a resting period, did not include participants who had suffered a sport-related concussion or did not present original research. Online supplementary table S1 provides a summary of the included studies.

A retrospective review of 184 athletes found that those individuals who were recommended to have a period of cognitive rest had a longer duration of symptoms than those who did not.<sup>7</sup> However, the individuals who were prescribed rest may have been perceived to have suffered a more severe injury. The effect of prescribed rest on neurocognitive test performance and symptom reporting was evaluated in high school and collegiate athletes who presented to an outpatient clinic at varying times since injury.<sup>8</sup> Regardless of the time since injury (eg, less than 1 week or greater than a month), the athlete was prescribed 1 week of complete physical and cognitive rest. Significant improvements in Post-Concussion Symptom Scale ratings and ImpACT composite scores were evident following 1 week of rest. However, there was no control or comparison group, so the improvements could have been attributed to a diverse range of factors.

Majerske *et al*<sup>9</sup> followed a group of 95 student-athletes presenting to a university hospital sport medicine concussion programme following a sport-related concussion. The athletes were retrospectively assigned to groups based on their self-reported activity level at the time of their visits (see online supplementary table S1). Those who reported moderate levels of cognitive and physical exertion (eg, participation in school and light activity at home such as jogging) over the first month after injury appeared to have better neuropsychological outcomes than those with very little (eg, no school or exercise) or high levels of activity (eg, school activity and participation in a sports game). Individuals reporting a high level of activity performed the worst on measures of visual memory and reaction time. Although causal explanations cannot be drawn from this observational study, their findings suggest that under-activity and over-activity might both be counterproductive for some injured athletes.

### Treatment

Twelve studies met the inclusion criteria evaluating the effect of treatment. The majority of abstracts reviewed that did not meet the inclusion criteria were not original research or did not evaluate the effect of treatment in individuals who had suffered a sport-related concussion. The nature of the treatment and the patients varied enough that a meta-analysis was not possible.

### Pharmacotherapy

Medications have been prescribed in case studies of athletes with headaches and neuroendocrine problems. One group pharmacological study has been published. Treatment with amantadine (100 mg twice daily for 3–4 weeks) was evaluated in individuals with symptoms greater than 21 days.<sup>5</sup> Data were collected retrospectively in a cohort of individuals who received amantadine as treatment and compared with historical controls (who were treated at the same clinic prior to the clinic utilising amantadine as treatment). All patients improved over time for

the measures taken, but the amantadine group improved more on measures of verbal memory, reaction time and number of symptoms endorsed. However, the individuals in the treatment group were at a lower baseline point at the start of the study for verbal memory and visual memory, and they reported more symptoms. Also, there may have been other differences between the groups (aside from amantadine) that could have accounted for any observed effect.

### Exercise

Three articles that met the inclusion criteria evaluated the effects of exercise as treatment, including one case study.<sup>10–12</sup> The effect of a combined treatment approach was evaluated in a case series of 16 children and adolescents (11 boys and 5 girls; median age 14, range 11–18 years) with persistent symptoms following concussion.<sup>10</sup> The treatment regimen consisted of a combined approach of submaximal aerobic training, light sport-specific coordination exercises, visualisation and imagery techniques, and a home exercise programme. The mean duration of treatment was 4.4 weeks (95% CI 3.1 to 5.7), and all children were able to resume normal physical activity participation following the treatment period.<sup>10</sup>

The effect of the subsymptom threshold daily aerobic exercise programme was evaluated in a crossover study of 12 patients with persistent symptoms following a concussion.<sup>12</sup> Symptoms were present at rest for a duration greater than 6 weeks but less than 1 year. All patients had symptoms during a graded exercise treadmill test. They completed a treadmill exercise test before and after a 2 or 3 week baseline non-intervention period. All patients continued onto the treatment phase at the completion of the baseline phase. Participants exercised 5–6 times per week until they were asymptomatic with exhaustive exercise. An overall reduction was found in the mean symptom score between baseline and treatment periods, and all patients were able to return to preinjury levels of activity.<sup>12</sup>

Participating in a 1 week regimen of hyperbaric oxygen therapy was associated with improvements in mental activation and cognitive flexibility in a case crossover design compared to a 1 week period of rest.<sup>13</sup> A case series of elite athletes with persistent symptoms of dizziness, neck pain and headaches had functional improvements and reported symptomatic improvements following a course of combined cervical spine manual therapy, neuromotor retraining, sensorimotor retraining and vestibular physiotherapy treatment following a sport-related concussion.<sup>14</sup> A randomised controlled trial including individuals with persistent symptoms of dizziness, neck pain and headaches following a sport-related concussion found that individuals treated with a combination of cervical spine manual therapy, neuromotor retraining, sensorimotor retraining and vestibular physiotherapy treatment were more likely to be medically cleared to return to sport within 8 weeks of initiating treatment (risk ratio 10.3; 95% CI 1.51 to 69.6).<sup>14 15</sup>

### DISCUSSION

There are very few published studies evaluating the effects of various forms of treatment or rest in athletes who have suffered a sport-related concussion. The included studies were case studies<sup>6 11 16 17</sup> (n=4) or case series (n=6),<sup>4 8–10 14 18</sup> quasi-experimental (n=1),<sup>5</sup> case crossover (n=2)<sup>12 13</sup> and one randomised controlled trial.<sup>15</sup> The overall quality of the literature reviewed lacked randomisation and many designs were retrospective and did not include a comparison group. In order to provide a more comprehensive discussion in this area,

additional references have been brought into the discussion from the non-sport-related concussion literature where appropriate.

### How much rest is too much?

In the initial days following a concussion, mental and physical rest has been strongly encouraged.<sup>19 20</sup> Three lines of evidence indirectly support the value of rest.<sup>21 22</sup> First, concussions can have a large adverse effect on physical and cognitive functioning in the first few days postinjury.<sup>23</sup> Acutely, the brain might be in a state of neurometabolic crisis,<sup>24 25</sup> at which time increased energy demand may hinder the restorative process, and it is believed that rest might facilitate recovery. Second, in animal injury models, there appears to be a 'temporal window' of vulnerability in which a second overlapping injury results in greater levels of traumatic axonal injury and magnified cognitive and behavioural deficits.<sup>26–28</sup> Thus, a rest period will reduce the likelihood of the athlete experiencing an overlapping injury. Finally, it has been demonstrated in rodent models that exercise appears to be good for the injured brain; however, animals that are allowed to exercise too soon after injury do not show the expected exercise-induced increases in molecular markers of neuroplasticity.<sup>29–32</sup> For these reasons, it is believed that rest is very likely beneficial following injury. However, this is largely based on animal research, theory and expert consensus.

At present, there are no evidence-based guidelines for how to manage athletes with slow recovery. The optimal time period for rest is unknown. Moreover, the specific schedule and type of rest (eg, bed rest versus greatly restricted activities) have not been studied and it has recently been argued that absolute rest is unrealistic.<sup>33</sup> In general, prolonged bed rest following medical procedures or as an intervention for health problems has rarely been shown to be beneficial and may cause harm.<sup>34</sup> There has been only one clinical trial involving bed rest following mild traumatic brain injury (MTBI) in civilians recruited from the emergency department, and this trial did not support the use of 6 days of bed rest as a management strategy.<sup>35</sup>

Although resting until symptom free is widely recommended following a concussion, only three studies met the inclusion criteria evaluating rest. One case crossover study did not show improvements in symptoms during a period of non-intervention,<sup>12</sup> and a case series demonstrated a longer duration of symptoms in individuals who were prescribed cognitive rest.<sup>7</sup> Positive effects on neurocognitive function and symptoms were reported following a 1-week period of prescribed rest in another study irrespective of the time since injury.<sup>8</sup> However, no long-term follow-up was discussed, so the impact that the resting period had on the ability to resume activity is not known. Additionally, individuals were not randomised to rest versus other activity levels, so it is not known if improvements in cognitive function and symptom reports were due to physical rest, cognitive rest, combined rest, non-specific effects of contact with a specialist, education and reassurance or other factors. Further studies to evaluate the effects of a resting period and the optimal duration of this period are needed. In the absence of evidence-based recommendations, a sensible approach involves the gradual return to school and social activities (prior to contact sports) in a manner that does not result in a significant exacerbation of symptoms.

### Pharmacotherapy

A wide variety of medications, such as non-steroidal anti-inflammatories, sleep agents, methylphenidate, antidepressants and anticonvulsants, have been used for persistent headache,

sleep problems and other symptoms following MTBI in civilians and military personnel. Clinical practice guidelines have been developed for how to manage civilians, active duty military personnel and veterans who are slow to recover from MTBI.<sup>36</sup> However, there is very little research evaluating pharmacological treatment following sport-related concussion. There is evidence that cases with a secondary diagnoses may respond to medical intervention. One study suggests that amantadine may facilitate the alleviation of symptoms and promote neurocognitive recovery in some athletes. However, although greater improvements were reported in the medication group, the control group and all patients demonstrated improvement over time.<sup>5</sup>

### Using exercise as treatment for athletes who are slow to recover

From a practical perspective, athletes need to, and naturally will, transition back into an active lifestyle. If their normal activities are restricted for extended periods of time, they are at risk for secondary problems such as physical deconditioning, anxiety and stress, mild depression, and irritability. Over time, the strength of the relation between the neurobiology of the injury and the ongoing symptoms very likely diminishes, and the pre-existing, current and contextual factors increasingly contribute to causing, maintaining or exacerbating symptoms. In the absence of good scientific evidence, clinicians must decide when to transition from activity restrictions and watchful waiting to more active treatment and rehabilitation, including exercise.

Converging lines of diverse, albeit indirect, medical and scientific evidence support the use of exercise as a core component of treatment for children, adolescents and young adults who are slow to recover from concussion.<sup>21 22</sup> Some of the benefits of exercise that have been reported include facilitation of molecular markers of neuroplasticity and neurogenesis,<sup>29 30 37–41</sup> improved cognitive functioning,<sup>42–45</sup> greater bilateral hippocampal volumes and better performance on a memory test in children with greater fitness levels,<sup>46 47</sup> changes in neurotransmitter systems,<sup>48 49</sup> higher ratings of self-esteem<sup>50</sup> and improved sleep quality.<sup>51–53</sup> Exercise has also been reported to be an effective adjunctive treatment in adults for depression<sup>54–61</sup> and anxiety,<sup>62–67</sup> chronic fatigue<sup>68 69</sup> and migraines.<sup>70 71</sup> Interestingly, adults who exercise regularly, and then abruptly stop for a few days (and up to 2 weeks), report increased levels of fatigue, a negative mood and symptoms of depression.<sup>72–75</sup> Thus, there may be non-specific adverse effects of abrupt discontinuation of exercise in concussed athletes that mimic and/or magnify ongoing symptoms of concussion.

Two published studies have examined exercise as a component of active rehabilitation for children, adolescents or adults who are slow to recover from a concussion.<sup>14 20</sup> Symptoms were reported to decrease and functional levels were reported to increase in these studies. However, further studies evaluating the effect of exercise compared to rest are needed because none of the currently available published studies included randomisation of patients to the treatment group, making a causal association difficult to determine and allowing for the potential of selection bias.

### Psychological intervention

Researchers have successfully used psychological treatment, such as modified forms of cognitive behavioural therapy, with adults who have chronic symptoms and problems following MTBI.<sup>76–79</sup> Psychological treatment is recommended in clinical practice guidelines for MTBI in civilians and the military,<sup>36</sup> and it has been strongly promoted as a component of a sequenced

care model for MTBI.<sup>80</sup> There is a large and mature literature indicating that psychological treatments are effective for reducing symptoms and improving functioning in adults with depression<sup>81–82</sup> and generalised anxiety disorder.<sup>83</sup> Behavioural and psychological treatments are effective for improving sleep and reducing psychological distress in people with insomnia.<sup>84</sup> Therefore, although not studied in sport-related concussion, it is plausible that psychological treatment for athletes who are slow to recover may be of benefit and should be studied more systematically.

### Rehabilitative treatment

There is some evidence that vestibular rehabilitation is effective for dizziness and balance problems in children and adults who have sustained an MTBI.<sup>85–86</sup> Commonly reported symptoms following a concussion include headaches, dizziness, nausea and neck pain.<sup>87</sup> Rehabilitation aimed at reducing symptoms and facilitating functional recovery of balance may be of benefit, as illustrated in one paper included in this review.<sup>10</sup> A randomised controlled trial and a case series reported benefit associated with a combined approach of orthopaedic and vestibular physiotherapy treatment following a concussion.<sup>14–15</sup> It may be that the cervical spine and/or balance centres are injured in some athletes and may benefit from treatment targeting these areas.<sup>88</sup> If the assessment of a patient with persistent headache after a concussion suggests that there is a cervicogenic aetiology, then cervical spine manual therapy and specific neuromotor control exercises for the cervicospinal region have been demonstrated to have benefit in relieving the headache.<sup>89–90</sup>

### CONCLUSIONS

- ▶ The current evidence evaluating the effect of rest and treatment following a sport-related concussion is sparse.
- ▶ An initial period of rest may be of benefit. However, further research to evaluate the long-term outcome of rest, and the optimal amount and type of rest, is needed.
- ▶ Low-level exercise may be of benefit, although the optimal timing postinjury for initiation of this treatment is currently unknown.
- ▶ Rehabilitative techniques in individuals with clinical findings suggesting cervical spine or vestibular dysfunction may facilitate recovery.
- ▶ There is a strong need for high-level studies evaluating the effects of a resting period, pharmacological interventions, rehabilitative techniques and exercise for individuals who are slow to recover from a sport-related concussion.

**Acknowledgements** The authors thank Chantel Debert for her review of the manuscript.

**Contributors** All authors participated in the initial formation of the questions and search strategies. KJS performed the searches, screening of titles and abstracts, critical appraisal of studies, writing of the manuscript and extraction of the data. GLI wrote portions of the manuscript and assisted with data extraction and searches. CAE reviewed questionable abstracts and assisted with data extraction and writing of the manuscript. SAH and PM assisted with writing of the manuscript and review of the included studies. WHM reviewed questionable abstracts and assisted with data extraction and writing of the manuscript.

**Competing interests** See the supplementary online data for competing interests (<http://dx.doi.org/10.1136/bjsports-2013-092190>).

**Provenance and peer review** Commissioned; internally peer reviewed.

### REFERENCES

- 1 McCrory P, Meeuwisse W, Johnston K, *et al*. Consensus Statement on Concussion in Sport 3rd International Conference on Concussion in Sport Held in Zurich, November 2008. *Clin J Sport Med* 2009;19:185–200.
- 2 McCrory PJK, Meeuwisse W, Aubry M, *et al*. Summary and agreement statement of the 2nd International Conference on Concussion in Sport, Prague 2004. *Br J Sports Med* 2005;39:196–204.
- 3 Group OCFEBMLoEW. The Oxford 2011 levels of evidence. *Oxf Centre Evid Based Med* 2011 (accessed 18 Sep 2012).
- 4 Tanriverdi F, Unluhizarci K, Kelestimur F. Pituitary function in subjects with mild traumatic brain injury: a review of literature and proposal of a screening strategy. *Pituitary* 2010;13:146–53.
- 5 Reddy CC, Collins M, Lovell M, *et al*. Efficacy of amantadine treatment on symptoms and neurocognitive performance among adolescents following sports-related concussion. *J Head Trauma Rehabil* 2012.
- 6 Foley CM, Wang DH. Central diabetes insipidus following a sports-related concussion: a case report. *Sports Health* 2012;4:139–41.
- 7 Gibson S, Meehan W. The effect of cognitive rest on duration of sport-related concussion symptoms. *Med Sci Sports Exerc* 2010;42:752.
- 8 Moser RS, Glatts C, Schatz P. Efficacy of immediate and delayed cognitive and physical rest for treatment of sports-related concussion. *J Pediatr* 2012; 161:922–6.
- 9 Majerske C, Mihalik J, Ren D, *et al*. Concussion in sports: postconcussive activity levels, symptoms and neurocognitive performance. *J Athl Train* 2008;43:265–74.
- 10 Gagnon I, Galli C, Friedman D, *et al*. Active rehabilitation for children who are slow to recover following sport-related concussion. *Brain Inj* 2009;23:956–64.
- 11 Zafonte R. Diagnosis and management of sports-related concussion: a 15-year-old athlete with a concussion. *JAMA* 2011;306:79–86.
- 12 Leddy JJ, Kozlowski K, Donnelly JP, *et al*. A preliminary study of subsymptom threshold exercise training for refractory post-concussion syndrome. *Clin J Sport Med* 2010;20:21–7.
- 13 Hardy P, Johnston K, Montgomery D, *et al*. Cognitive effects of hyperbaric oxygen therapy on sports-related concussion. *Clin J Sport Med* 2003;13:378–80.
- 14 Schneider K, Meeuwisse W, Emery C. Symptom and functional improvements following a course of vestibular rehabilitation, manual therapy and spinal stabilization exercise in high performance athletes with complex concussions. *Clin J Sport Med* 2009;19:265–6.
- 15 Schneider K, Meeuwisse W, Boyd L, *et al*. Multimodal physiotherapy in the treatment of individuals with persistent symptoms following a sport related concussion: a randomized controlled trial. *Clin J Sport Med* 2012;22:295.
- 16 Abend N, Nance M, Bonneman C. Subcutaneous sumatriptan in an adolescent with acute posttraumatic headaches. *J Child Neurol* 2008;23:438–40.
- 17 Ives J, Alderman M, Stred S. Hypopituitarism after multiple concussions: a retrospective case study in an adolescent male. *J Athl Train* 2007;42:431–9.
- 18 Guskiewicz KM, Weaver NL, Padua DA, *et al*. Epidemiology of concussion in collegiate and high school football players. *Am J Sports Med* 2000;28:643–50.
- 19 McCrory P, Johnston K, Aubry M, *et al*. Summary of the second international conference on concussion in sport, Prague, Czech Republic. *Clin J Sport Med* 2005;15:48–55.
- 20 McCrory P, Meeuwisse W, Johnston K, *et al*. Consensus statement on concussion in sport: the 3rd International Conference on Concussion in Sport held in Zurich, November 2008. *Br J Sports Med* 2009;43(Suppl 1):i76–90.
- 21 Silverberg ND, Iverson GL. Is rest after concussion 'The Best Medicine?': recommendations for activity resumption following concussion in athletes, civilians, and military service members. *J Head Trauma Rehabil* 2012.
- 22 Iverson GL, Gagnon I, Griesbach GS. Active rehabilitation for slow-to-recover children following mild traumatic brain injury. In: Yeates KO, Kirkwood M. eds. *Mild traumatic brain injury in children and adolescents: from basic science to clinical management*. New York: Guilford Press, 2012:281–302.
- 23 Broglio SP, Puetz TW. The effect of sport concussion on neurocognitive function, self-report symptoms and postural control: a meta-analysis. *Sports Med* 2008;38:53–67.
- 24 Giza CC, Hovda DA. The pathophysiology of traumatic brain injury. In: Lovell MR, Echemendia RJ, Barth JT, Collins MW. eds. *Traumatic brain injury in sports*. Lisse: Swets & Zeitlinger, 2004:45–70.
- 25 Giza CC, Hovda DA. The neurometabolic cascade of concussion. *J Athl Train* 2001;36:228–35.
- 26 Vagnozzi R, Tavazzi B, Signoretti S, *et al*. Temporal window of metabolic brain vulnerability to concussions: mitochondrial-related impairment—part I. *Neurosurgery* 2007;61:379–88; discussion 388–9.
- 27 Longhi L, Saatman KE, Fujimoto S, *et al*. Temporal window of vulnerability to repetitive experimental concussive brain injury. *Neurosurgery* 2005;56:364–74; discussion 364–74.
- 28 Laurer HL, Bareyre FM, Lee VM, *et al*. Mild head injury increasing the brain's vulnerability to a second concussive impact. *J Neurosurg* 2001;95:859–70.
- 29 Griesbach GS, Hovda DA, Molteni R, *et al*. Voluntary exercise following traumatic brain injury: brain-derived neurotrophic factor upregulation and recovery of function. *Neuroscience* 2004;125:129–39.
- 30 Griesbach GS, Sutton RL, Hovda DA, *et al*. Controlled contusion injury alters molecular systems associated with cognitive performance. *J Neurosci Res* 2009;87:795–805.

- 31 Griesbach GS, Gomez-Pinilla F, Hovda DA. The upregulation of plasticity-related proteins following TBI is disrupted with acute voluntary exercise. *Brain Res* 2004;1016:154–62.
- 32 Griesbach GS. Exercise after traumatic brain injury: is it a double-edged sword? *PM R* 2011;3(6 Suppl 1):S64–72.
- 33 Kirkwood M, Randolph C, Yeates K. Sport-related concussion: a call for evidence and perspective amidst the alarms. *Clin J Sport Med* 2012;22:383–4.
- 34 Allen C, Glasziou P, Del Mar C. Bed rest: a potentially harmful treatment needing more careful evaluation. *Lancet* 1999;354:1229–33.
- 35 de Kruijk JR, Leffers P, Meerhoff S, et al. Effectiveness of bed rest after mild traumatic brain injury: a randomised trial of no versus six days of bed rest. *J Neurol Neurosurg Psychiatry* 2002;73:167–72.
- 36 Group MocomW. VA/DoD Clinical Practice Guideline for management of concussion/mild traumatic brain injury. *J Rehabil Res Dev* 2009;46:CP1–68.
- 37 Michelini LC, Stern JE. Exercise-induced neuronal plasticity in central autonomic networks: role in cardiovascular control. *Exp Physiol* 2009;94:947–60.
- 38 Neeper SA, Gomez-Pinilla F, Choi J, et al. Exercise and brain neurotrophins. *Nature* 1995;373:109.
- 39 van Praag H. Neurogenesis and exercise: past and future directions. *Neuromolecular Med* 2008;10:128–40.
- 40 Griesbach GS, Gomez-Pinilla F, Hovda DA. Time window for voluntary exercise-induced increases in hippocampal neuroplasticity molecules after traumatic brain injury is severity dependent. *J Neurotrauma* 2007;24:1161–71.
- 41 Griesbach GS, Hovda DA, Gomez-Pinilla F, et al. Voluntary exercise or amphetamine treatment, but not the combination, increases hippocampal brain-derived neurotrophic factor and synapsin I following cortical contusion injury in rats. *Neuroscience* 2008;154:530–40.
- 42 Smith PJ, Blumenthal JA, Hoffman BM, et al. Aerobic exercise and neurocognitive performance: a meta-analytic review of randomized controlled trials. *Psychosom Med* 2010;72:239–52.
- 43 Castelli DM, Hillman CH, Buck SM, et al. Physical fitness and academic achievement in third- and fifth-grade students. *J Sport Exerc Psychol* 2007;29:239–52.
- 44 Chaddock L, Hillman CH, Buck SM, et al. Aerobic fitness and executive control of relational memory in preadolescent children. *Med Sci Sports Exerc* 2011;43:344–9.
- 45 Buck SM, Hillman CH, Castelli DM. The relation of aerobic fitness to stroop task performance in preadolescent children. *Med Sci Sports Exerc* 2008;40:166–72.
- 46 Chaddock L, Erickson KI, Prakash RS, et al. A neuroimaging investigation of the association between aerobic fitness, hippocampal volume, and memory performance in preadolescent children. *Brain Res* 2010;1358:172–83.
- 47 Chaddock L, Erickson KI, Prakash RS, et al. Basal ganglia volume is associated with aerobic fitness in preadolescent children. *Dev Neurosci* 2010;32:249–56.
- 48 Molteni R, Ying Z, Gomez-Pinilla F. Differential effects of acute and chronic exercise on plasticity-related genes in the rat hippocampus revealed by microarray. *Eur J Neurosci* 2002;16:1107–16.
- 49 Chaouloff F. Physical exercise and brain monoamines: a review. *Acta Physiol Scand* 1989;137:1–13.
- 50 Ekeland E, Heian F, Hagen KB, et al. Exercise to improve self-esteem in children and young people. *Cochrane Database Syst Rev* 2004;1:CD003683.
- 51 Youngstedt SD. Effects of exercise on sleep. *Clin Sports Med* 2005;24:355–65.
- 52 Brand S, Beck J, Gerber M, et al. 'Football is good for your sleep': favorable sleep patterns and psychological functioning of adolescent male intense football players compared to controls. *J Health Psychol* 2009;14:1144–55.
- 53 Brand S, Gerber M, Beck J, et al. High exercise levels are related to favorable sleep patterns and psychological functioning in adolescents: a comparison of athletes and controls. *J Adolesc Health* 2010;46:133–41.
- 54 Penninx BW, Rejeski WJ, Pandya J, et al. Exercise and depressive symptoms: a comparison of aerobic and resistance exercise effects on emotional and physical function in older persons with high and low depressive symptomatology. *J Gerontol B Psychol Sci Soc Sci* 2002;57:124–32.
- 55 Dunn AL, Trivedi MH, Kampert JB, et al. Exercise treatment for depression: efficacy and dose response. *Am J Prev Med* 2005;28:1–8.
- 56 Mead GE, Morley W, Campbell P, et al. Exercise for depression. *Cochrane Database Syst Rev* 2008(4):CD004366.
- 57 Babyak M, Blumenthal JA, Herman S, et al. Exercise treatment for major depression: maintenance of therapeutic benefit at 10 months. *Psychosom Med* 2000;62:633–8.
- 58 Lawlor DA, Hopker SW. The effectiveness of exercise as an intervention in the management of depression: systematic review and meta-regression analysis of randomised controlled trials. *BMJ* 2001;322:763–7.
- 59 Daley A. Exercise and depression: a review of reviews. *J Clin Psychol Med Set* 2008;15:140–7.
- 60 Mead GE, Morley W, Campbell P, et al. Exercise for depression. *Cochrane Database Syst Rev* 2009(3):CD004366.
- 61 Rethorst CD, Wipfli BM, Landers DM. The antidepressive effects of exercise: a meta-analysis of randomized trials. *Sports Med* 2009;39:491–511.
- 62 Wang C, Bannuru R, Ramel J, et al. Tai Chi on psychological well-being: systematic review and meta-analysis. *BMC Complement Altern Med* 2010;10:23.
- 63 Herring MP, O'Connor PJ, Dishman RK. The effect of exercise training on anxiety symptoms among patients: a systematic review. *Arch Intern Med* 2010;170:321–31.
- 64 Greenwood BN, Fleshner M. Exercise, learned helplessness, and the stress-resistant brain. *Neuromolecular Med* 2008;10:81–98.
- 65 Barbour KA, Edenfield TM, Blumenthal JA. Exercise as a treatment for depression and other psychiatric disorders: a review. *J Cardiopulm Rehabil Prev* 2007;27:359–67.
- 66 Merom D, Phongsavan P, Wagner R, et al. Promoting walking as an adjunct intervention to group cognitive behavioral therapy for anxiety disorders—a pilot group randomized trial. *J Anxiety Disord* 2008;22:959–68.
- 67 Smits JA, Berry AC, Rosenfield D, et al. Reducing anxiety sensitivity with exercise. *Depress Anxiety* 2008;25:689–99.
- 68 Edmonds M, McGuire H, Price J. Exercise therapy for chronic fatigue syndrome. *Cochrane Database Syst Rev* 2004(3):CD003200.
- 69 White PD, Goldsmith KA, Johnson AL, et al. Comparison of adaptive pacing therapy, cognitive behaviour therapy, graded exercise therapy, and specialist medical care for chronic fatigue syndrome (PACE): a randomised trial. *Lancet* 2011;377:823–36.
- 70 Lockett DM, Campbell JF. The effects of aerobic exercise on migraine. *Headache* 1992;32:50–4.
- 71 Koseoglu E, Akboyraz A, Soyuer A, et al. Aerobic exercise and plasma beta endorphin levels in patients with migrainous headache without aura. *Cephalalgia* 2003;23:972–6.
- 72 Weinstein AA, Deuster PA, Kop WJ. Heart rate variability as a predictor of negative mood symptoms induced by exercise withdrawal. *Med Sci Sports Exerc* 2007;39:735–41.
- 73 Berlin AA, Kop WJ, Deuster PA. Depressive mood symptoms and fatigue after exercise withdrawal: the potential role of decreased fitness. *Psychosom Med* 2006;68:224–30.
- 74 Glass JM, Lyden AK, Petzke F, et al. The effect of brief exercise cessation on pain, fatigue, and mood symptom development in healthy, fit individuals. *J Psychosom Res* 2004;57:391–8.
- 75 Mondin GW, Morgan WP, Piering PN, et al. Psychological consequences of exercise deprivation in habitual exercisers. *Med Sci Sports Exerc* 1996;28:1199–203.
- 76 Potter S, Brown RG. Cognitive behavioural therapy and persistent post-concussional symptoms: integrating conceptual issues and practical aspects in treatment. *Neuropsychol Rehabil* 2012;22:1–25.
- 77 Ferguson RJ, Mittenberg W. Cognitive-behavioral treatment of postconcussion syndrome: a therapist's manual. In: Van Hasselt VB, Hersen M, eds. *Sourcebook of psychological treatment manuals for adult disorders*. New York: Plenum, 1996:615–55.
- 78 Miller LJ, Mittenberg W. Brief cognitive behavioral interventions in mild traumatic brain injury. *Appl Neuropsychol* 1998;5:172–83.
- 79 Leonard KN, Tucker DM. Group-based cognitive-behavioral therapy for persistent postconcussion syndrome: a controlled treatment outcome study. Paper presented at: International Neuropsychological Society; February, 2004; Baltimore, MD.
- 80 Iverson GL, Silverberg ND, Lange RT, et al. Conceptualizing outcome from mild traumatic brain injury. In: Zasler ND, Katz DI, Zafonte R, eds. *Brain injury medicine: principles and practice*, 2nd edn. New York: Demos Medical Publishing, 2012:470–97.
- 81 Bortolotti B, Menchetti M, Bellini F, et al. Psychological interventions for major depression in primary care: a meta-analytic review of randomized controlled trials. *Gen Hosp Psychiatry* 2008;30:293–302.
- 82 Ekers D, Richards D, Gilbody S. A meta-analysis of randomized trials of behavioural treatment of depression. *Psychol Med* 2008;38:611–23.
- 83 Hunot V, Churchill R, Silva de Lima M, et al. Psychological therapies for generalised anxiety disorder. *Cochrane Database Syst Rev*. 2007(1):CD001848.
- 84 Zeitzer JM, Hubbard J, Litsch S, et al. *Sleep Disorders in the context of traumatic brain injury. State of the Art (SOTA)*. Arlington, VA: Department of Veterans Affairs, 2008.
- 85 Alsalameh BA, Mucha A, Morris LO, et al. Vestibular rehabilitation for dizziness and balance disorders after concussion. *J Neurol Phys Ther* 2010;34:87–93.
- 86 Gottshall K. Vestibular rehabilitation after mild traumatic brain injury with vestibular pathology. *NeuroRehabilitation* 2011;29:167–71.
- 87 Benson BW, Meeuwisse WH, Rizos J. A prospective study of concussions among National Hockey League players during regular season games: the NHL-NHLPA Concussion Program. *CMAJ* 2011;183:905–11.
- 88 Schneider K, Meeuwisse W, Nettel-Aguirre A, et al. Cervico-vestibular physiotherapy in the treatment of individuals with persistent symptoms following sport related concussion: a randomized controlled trial (abstract). *Clin J Sport Med* 2012;22:295.
- 89 Lucas S, Hoffman JM, Bell KB, et al. Characterization of headache after traumatic brain injury. *Cephalalgia* 2012;32:600–6.
- 90 Jull G, Trott P, Potter H, et al. A randomized controlled trial of exercise and manipulative therapy for cervicogenic headache. *Spine* 2002;27:1835–43.



## The effects of rest and treatment following sport-related concussion: a systematic review of the literature

Kathryn J Schneider, Grant L Iverson, Carolyn A Emery, et al.

*Br J Sports Med* 2013 47: 304-307

doi: 10.1136/bjsports-2013-092190

---

Updated information and services can be found at:  
<http://bjsm.bmj.com/content/47/5/304.full.html>

---

*These include:*

**Data Supplement**

"Supplementary Data"

<http://bjsm.bmj.com/content/suppl/2013/03/11/47.5.304.DC1.html>

**References**

This article cites 76 articles, 14 of which can be accessed free at:

<http://bjsm.bmj.com/content/47/5/304.full.html#ref-list-1>

**Email alerting service**

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

**Topic Collections**

Articles on similar topics can be found in the following collections

[Physiotherapy](#) (88 articles)  
[Physiotherapy](#) (135 articles)

---

**Notes**

---

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>